Bentley University Tuberculosis (TB) Medical Provider Form

[This form is only required if the TB Risk Assessment had a "yes" response]

Name: _____
Last First M.I.

Date of Birth: ____ /____ /_____ Country of Birth: _____

Home Address: _____

Phone Number: (Home)_____ (Cell) _____

Step 1: Tuberculosis Testing – IGRA is preferred, but TST is also acceptable. Must be performed no earlier than 12 months prior to semester start, but preferably within 6 months prior.

If this test has been previously done and resulted "positive," skip to Step 2.

Date of IGRA (QuantiFERON Gold or T-Spot Blood Test): ___ /____ /____ **Result:** _____

*Please attach copy of actual lab results from the laboratory testing facility OR provider signed documentation

OR

Tuberculin Skin Test (TST) *(not recommended for those who have received BCG vaccine)*

Plant date: __ /__ /____ **Read date*:** __ /__ /____ **Result**:____mm of induration Positive / Negative

*48-72 hours after plant date ** See interpretation guidelines below

Step 2: If Positive IGRA or TST testing in Step 1, a Chest X-ray is required:

Date of Chest X-ray: ___ /___ /_____ **Result (Circle one):** Normal Abnormal

Clinical Evaluation: _____Normal _____Abnormal

Describe: _____

Treatment: _____No _____Yes

If Yes, list drug(s), dose frequency, and dates: _____

Health Care Provider

Name: _____ **Address:** _____

Signature: _____ **Date:** _____ **Phone:** _____

**** TST Interpretation Guidelines:**

5 mm or greater is positive:

- Recent close contacts of an individual with infectious TB
- Persons with fibrotic changes on a prior chest x-ray consistent with past TB disease
- Organ transplant recipients and other immunosuppressed persons (including receiving equivalent of > 15 mg/d of prednisone for > 1 month)
- Persons with HIV/AIDS

10 mm or greater is positive:

- Persons born in a high prevalence country or who resided in one for a significant amount of time
- Injection drug users
- Mycobacteriology laboratory personnel
- History of resident, worker, or volunteer in high-risk congregate settings
- Persons with the following clinical conditions: silicosis, diabetes mellitus chronic renal failure, certain types of cancer (leukemias and lymphomas, cancers of the head, neck, or lung), gastrectomy or intestinal bypass, chronic malabsorption syndromes, and low body weight (>10% below ideal)

15 mm or greater is positive:

- Persons with no known risk factors for TB disease